

10 tips

Nutrition
Education Series

salt and sodium

10 tips to help you cut back



It's clear that Americans have a taste for salt, but salt plays a role in high blood pressure. Everyone, including kids, should reduce their sodium intake to less than 2,300 milligrams of sodium a day (about 1 teaspoon of salt). Adults age 51 and older, African Americans of any age, and individuals with high blood pressure, diabetes, or chronic kidney disease should further reduce their sodium intake to 1,500 mg a day.

1 think fresh

Most of the sodium Americans eat is found in processed foods. Eat highly processed foods less often and in smaller portions—especially cheesy foods, such as pizza; cured meats, such as bacon, sausage, hot dogs, and deli/luncheon meats; and ready-to-eat foods, like canned chili, ravioli, and soups. Fresh foods are generally lower in sodium.

2 enjoy home-prepared foods

Cook more often at home—where you are in control of what's in your food. Preparing your own foods allows you to limit the amount of salt in them.

3 fill up on veggies and fruits—they are naturally low in sodium

Eat plenty of vegetables and fruits—fresh or frozen. Eat a vegetable or fruit at every meal.

4 choose dairy and protein foods that are lower in sodium

Choose more fat-free or low-fat milk and yogurt in place of cheese, which is higher in sodium. Choose fresh beef, pork, poultry, and seafood, rather than those with salt added. Deli or luncheon meats, sausages, and canned products like corned beef are higher in sodium. Choose unsalted nuts and seeds.

5 adjust your taste buds

Cut back on salt little by little—and pay attention to the natural tastes of various foods. Your taste for salt will lessen over time.



6 skip the salt

Skip adding salt when cooking. Keep salt off the kitchen counter *and* the dinner table. Use spices, herbs, garlic, vinegar, or lemon juice to season foods or use no-salt seasoning mixes. Try black or red pepper, basil, curry, ginger, or rosemary.

7 read the label

Read the Nutrition Facts label and the ingredients statement to find packaged and canned foods lower in sodium. Look for foods labeled “low sodium,” “reduced sodium,” or “no salt added.”

8 ask for low-sodium foods when you eat out

Restaurants may prepare lower sodium foods at your request and will serve sauces and salad dressings on the side so you can use less.

9 pay attention to condiments

Foods like soy sauce, ketchup, pickles, olives, salad dressings, and seasoning packets are high in sodium. Choose low-sodium soy sauce and ketchup. Have a carrot or celery stick instead of olives or pickles. Use only a sprinkling of flavoring packets instead of the entire packet.

10 boost your potassium intake


Choose foods with potassium, which may help to lower your blood pressure. Potassium is found in vegetables and fruits, such as potatoes, beet greens, tomato juice and sauce, sweet potatoes, beans (white, lima, kidney), and bananas. Other sources of potassium include yogurt, clams, halibut, orange juice, and milk.

10 tips

Nutrition
Education Series

cut back on your kid's sweet treats

10 tips to decrease added sugars



Limit the amount of foods and beverages with added sugars your kids eat and drink. If you don't buy them, your kids won't get them very often. Sweet treats and sugary drinks have a lot of calories but few nutrients. Most added sugars come from sodas, sports drinks, energy drinks, juice drinks, cakes, cookies, ice cream, candy, and other desserts.

1 serve small portions

It's not necessary to get rid of all sweets and desserts. Show kids that a small amount of treats can go a long way. Use smaller bowls and plates for these foods. Have them share a candy bar or split a large cupcake.

2 sip smarter

Soda and other sweet drinks contain a lot of sugar and are high in calories. Offer water, 100% juice, or fat-free milk when kids are thirsty.



3 use the check-out lane that does not display candy

Most grocery stores will have a candy-free check-out lane to help moms out. Waiting in a store line makes it easy for children to ask for the candy that is right in front of their faces to tempt them.

4 choose not to offer sweets as rewards

By offering food as a reward for good behavior, children learn to think that some foods are better than other foods. Reward your child with kind words and comforting hugs, or give them non-food items, like stickers, to make them feel special.

5 make fruit the everyday dessert

Serve baked apples, pears, or enjoy a fruit salad. Or, serve yummy frozen juice bars (100% juice) instead of high-calorie desserts.



6 make food fun

Sugary foods that are marketed to kids are advertised as "fun foods." Make nutritious foods fun by preparing them with your child's help and being creative together. Create a smiley face with sliced bananas and raisins. Cut fruit into fun and easy shapes with cookie cutters.

7 encourage kids to invent new snacks

Make your own snack mixes from dry whole-grain cereal, dried fruit, and unsalted nuts or seeds. Provide the ingredients and allow kids to choose what they want in their "new" snack.



8 play detective in the cereal aisle

Show kids how to find the amount of total sugars in various cereals. Challenge them to compare cereals they like and select the one with the lowest amount of sugar.

9 make treats "treats," not everyday foods

Treats are great once in a while. Just don't make treat foods an everyday thing. Limit sweet treats to special occasions.

10 if kids don't eat their meal, they don't need sweet "extras"

Keep in mind that candy or cookies should not replace foods that are not eaten at meal time.



Center for Nutrition
Policy and Promotion

Go to www.ChooseMyPlate.gov for more information.

DG TipSheet No. 13

June 2011

USDA is an equal opportunity
provider and employer.